

## HEPATOPROTECTIVE ACTIVITY OF DIFFERENT LEAF EXTRACTS OF *TECOMARIA CAPENSIS* IN RATS

ELAMARAN TAMIL JOTHI\*<sup>1</sup>, V.RAVICHANDIRAN<sup>1</sup>, P. DURGA NITHYA<sup>2</sup>, V. SRIKANTH<sup>2</sup> & V.SUBA<sup>3</sup>

<sup>1</sup>Dept of Pharmacology, School of Pharmacy, Vels University, Palavaram, Chennai, T.N, <sup>2</sup>Dept of Pharmacology, Vignan Pharmacy College, Vadlamudi, Guntur DT, A.P, <sup>3</sup> Dept of Pharmacology, National Institute of Siddha, Thambaram, Chennai, T.N

Received: 14 Feb 2012, Revised and Accepted: 26 Mar 2012

### ABSTRACT

The present study was conducted to evaluate the hepatoprotective activity of ethyl acetate and ethanolic extracts of *Tecomaria capensis* against paracetamol and CCl<sub>4</sub> induced liver damage in rats. The ethyl acetate and ethanolic extracts of *Tecomaria capensis* were administered orally (in paracetamol induced method) and intraperitoneally (in CCl<sub>4</sub> induced method) to the animals with hepatotoxicity induced by paracetamol and CCl<sub>4</sub>. Silymarin was given as reference standard. All the test drugs were administered by suspending in DMSO solution. The plant extract was effective in protecting the liver against the injury induced by paracetamol and CCl<sub>4</sub> in rats. This was evident from significant reduction in serum enzymes alanine aminotransferase (ALT) or Serum Glutamate Pyruvate Transaminase (SGPT), aspartate aminotransferase (AST) or Serum Glutamate Oxaloacetate Transaminase (SGOT), alkaline phosphatase (ALP) and bilirubin. It was concluded from the result that the ethyl acetate and ethanolic extract of *Tecomaria capensis* possesses hepatoprotective activity against paracetamol and CCl<sub>4</sub> induced hepatotoxicity in rats.

**Keywords:** *Tecomaria capensis*, Paracetamol, CCl<sub>4</sub>, Hepatoprotective and hepatotoxicity.

### INTRODUCTION

Liver is one of the largest organs in human body and the chief site for intense metabolism and excretion. So it has a surprising role in the maintenance, performance and regulating homeostasis of the body. It is involved with almost all the biochemical pathways to growth, fight against disease, nutrient supply, energy provision and reproduction. The major functions of the liver are carbohydrate, protein and fat metabolism, detoxification, secretion of bile and storage of vitamin. Thus, to maintain a healthy liver is a crucial factor for overall health and well-being. But it is continuously and variedly exposed to environmental toxins, abused by poor drug habits, alcohol and prescribed and over-the-counter drug which can eventually lead to various liver ailments like hepatitis, cirrhosis and alcoholic liver disease. Thus liver diseases are some of the fatal diseases in the world today. They pose a serious challenge to international public health. Modern medicines have little to offer for alleviation of hepatic diseases and it is chiefly the plant based preparations which are employed for their treatment of liver disorders. But there are not much drugs available for the treatment of liver disorders.

Unfortunately, conventional or synthetic drugs used in the treatment of liver diseases are inadequate and sometimes can have serious side effects<sup>1</sup>. In the absence of a reliable liver protective drug in modern medicine, there are a number of medicinal preparations in Ayurveda recommended for the treatment of liver disorders<sup>2</sup>. In view of severe undesirable side effects of synthetic agents, there is growing focus to follow systematic research methodology and to evaluate scientific basis for the traditional herbal medicines that are claimed to possess hepatoprotective activity.

Therefore, many folk remedies from plant origin are tested for its potential antioxidant and hepatoprotective liver damage in experimental animal model. Carbon tetrachloride (CCl<sub>4</sub>) induced hepatotoxicity model is widely used for the study of hepatoprotective effects of drugs and plant extracts. *Tecomaria capensis* is an ever green plant in warm climate areas but loses its leaves in colder areas. It has pinnately compound leaves that have oval leaflets with blunt teeth and orange coloured flowers. Plant is used as traditional medicine to relieve pain and sleeplessness<sup>3</sup>. Dried powdered bark infusions are taken for sleeplessness<sup>4</sup>, reported to induce sleep<sup>5</sup>. It is included in the list of African plants evaluated for invitro antiplasmodial activity against *Plasmodium falciparum*<sup>6</sup>.

### MATERIALS AND METHODS

#### Plant materials and Preparation of Extracts

The leaves of *Tecomaria capensis* were collected from Guntur, Andhra Pradesh. It was authenticated by professor Dr.S.M.Khasim, Department of Botony and Microbiology, Acharya Nagarjuna University, Nagarjuna nagar, Guntur. The leaf part of *Tecomaria capensis* was dried at room temperature and grounded into powder and passed through 60# sieve. The powder (500gm) was extracted successively in soxhlet by ethanol and ethyl acetate. The sediments were filtered and the filtrate was dried at 400°C in an oven to get dried product. The different fractions obtained were used for further study.

#### Animals

Albino rats (150-250 gm. each) of either sex were kept under standard environmental conditions (25±2°C under 12 h light & 12 h dark cycles) in polypropylene cages. Standard pelleted feed & drinking water were provided ad libitum throughout the experimental period. The animals were acclimated to laboratory conditions, one week prior to the initiation of experimental work. The animals were divided into seven groups of six each.

#### Experimental design for hepatoprotective activity

##### CCl<sub>4</sub> induced method

The rats were divided randomly into seven groups with six rats in each. The hepatoprotective activity of the plant extracts was tested using CCl<sub>4</sub> model. Group I (control) received neither the plant extract nor CCl<sub>4</sub> for 21 days, Group II (negative control) was given a single intraperitoneal dose of CCl<sub>4</sub> (3ml/kg), Group III (positive control) was given intraperitoneal doses of CCl<sub>4</sub> (3ml/kg) and standard (Silymarin (25mg/kg), Group IV, V received crude ethanolic extract of plant material *Tecomaria capensis* at intraperitoneal dose of 100mg/kg b.wt. (Low dose) and 200mg/kg b.wt. (high dose) respectively and a single dose of CCl<sub>4</sub>, Group VI, VII received crude ethyl acetate extract of *Tecomaria capensis* at an intraperitoneal dose of 100mg/kg b.wt. (Low dose) and 200mg/kg b.wt. (High dose) respectively and a single dose of CCl<sub>4</sub><sup>7</sup>.

##### Paracetamol induced method

A total of 42 animals were equally divided into 7 groups of six each. Group - I served as control received neither the plant extract nor paracetamol, Group - II served as paracetamol control, administered with paracetamol (3gm/kg) as single dose, Group - III served as

reference control, received Silymarin (25mg/kg) once daily, Group - IV, V received *Tecomaria capensis* ethanolic extract low dose (100 mg/kg) and high dose (200 mg/kg) respectively once daily for 3 days, Group -VI, VII received *Tecomaria capensis* ethyl acetate extract low dose (100 mg/kg) and high dose (200 mg/kg) respectively once daily. Group III - VII received paracetamol (3gm/kg) as single dose thirty minutes after the administration of *Tecomaria capensis* extracts and silymarin<sup>8</sup>.

**Assessment of hepatoprotective activity**

Rats were treated as per the treatment protocol. Body weights of these rats were monitored sequentially in control and experimental animals for a period of 21 days for CCl<sub>4</sub> induced method and 3 days for paracetamol induced method.

**Biochemical estimation:** Rats were sacrificed 1 hour after administration on day 21. The blood was collected by retro orbital artery bleeding. Blood samples were centrifuged for 10 minutes at 3000 rpm to separate the serum. SGOT, SGPT, ALP and Total Bilirubin levels were estimated from the serum.

**Histopathological studies:** The livers were excised quickly and fixed in 10% formalin and stained with haemotoxylin and eosin and then observed under microscope for degeneration, fatty changes, necrotic changes and evidence of hepatotoxicity if any.

**Statistics:** The results were expressed as Mean ± SEM. Statistical analysis was carried out by using two tailed 't' test.

**RESULTS**

The results of hepatoprotective activities of crude ethylacetate and ethanolic extracts of this plant at a dose of 100 mg/kg b.wt. and 200mg/kg b.wt. on rats intoxicated with carbon tetrachloride and paracetamol were illustrated in the table 1 and table 2 respectively. The tables also showed the comparison of effects among the untreated (control) and carbon tetrachloride treated (negative

control) group with the drug treated group of rats. The results were represented as Mean ± Standard Error of Mean (M±SEM).

Carbon tetrachloride group significantly increased the serum level of SGOT (143.26%), SGPT (89.37%), Total Bilirubin (91.62%) and ALP (200.03%) as shown in table 1. The 't' value obtained exceeded the limit of significance. That is in all cases, p < 0.0001.

**Treatment Groups (Table 1):**

**Positive control (silymarin) Groups (G3):** There was significant decrease in total Bilirubin (0.7 ± 0.115 mg/dl), accompanied by significant decrease in level of SGOT (56.33 ± 2.33 mg/dl) and also significant decrease in ALP (38.83 ± 1.42 IU/dl), SGPT (75.50 ± 1.18 IU/dl), as compared to the negative control.

**Ethyl acetate low dose Groups (G4):** There was significant increase in total Bilirubin (1.667 ± 0.154 mg/dl), accompanied by significant decrease in level of SGOT (86.0 ± 3.35 mg/dl) and also significant decrease in ALP (76.17 ± 1.35 IU/dl), SGPT (108.67 ± 2.14 IU/dl), as compared to the negative control.

**Ethyl acetate high dose (G5):** There was significant decrease in total Bilirubin (1.2 ± 0.106 mg/dl), accompanied by significant decrease in level of SGOT (80.0 ± 2.0 mg/dl) and also significant decrease in ALP (55.67 ± 1.38 IU/dl), SGPT (94.83 ± 1.62 IU/dl), as compared to the negative control.

**Ethanol low dose (G6):** There was significant decrease in total Bilirubin (1.317 ± 0.158 mg/dl), accompanied by significant decrease in level of SGOT (75.50 ± 1.26 mg/dl) and also significant decrease in ALP (64.17 ± 1.74 IU/dl), SGPT (100.0 ± 1.67 IU/dl), as compared to the negative control.

**Ethanol high dose (G7):** There was significant decrease in total Bilirubin (1.117 ± 0.095 mg/dl), accompanied by significant decrease in level of SGOT (62.17 ± 0.83 mg/dl) and also significant decrease in ALP (44.83 ± 1.82 IU/dl), SGPT (82.33 ± 0.88 IU/dl), as compared to the negative control.

**Table 1: Effects of ethyl acetate and ethanolic extracts of *Tecomaria capensis* plant on various biochemical parameters in rats with carbon tetrachloride induced hepatotoxicity.**

| Group                  | SGOT        | SGPT         | ALP         | Total Bilirubin | Direct         | Indirect     |
|------------------------|-------------|--------------|-------------|-----------------|----------------|--------------|
| Control                | 49.33±1.73  | 67.50±1.43   | 28.33±1.71  | 0.8±0.115       | 0.3±0.058      | 0.417±0.095  |
| Negative control       | 120.0±2.73  | 127.83±1.25  | 85.0±1.06   | 1.533±0.115     | 0.7±0.152      | 0.8±0.106    |
| Positive control       | 56.33±2.33  | 75.50±1.18   | 38.83±1.42  | 0.7±0.115       | 0.4±0.106      | 0.4±0.073    |
| Ethylacetate low dose  | 86.0±3.35*  | 108.67±2.14* | 76.17±1.35* | 1.667±0.154**   | 0.6±0.097***   | 1.0±0.106**  |
| Ethylacetate high dose | 80.0±2.0*   | 94.83±1.62*  | 55.67±1.38* | 1.2±0.106***    | 0.717±0.154*** | 0.517±0.101  |
| Ethanol low dose       | 75.50±1.26* | 100.0±1.67*  | 64.17±1.74* | 1.317±0.158***  | 0.5±0.115      | 0.8±0.139*** |
| Ethanol high dose      | 62.17±0.83* | 82.33±0.88*  | 44.83±1.82* | 1.117±0.095     | 0.617±0.095*** | 0.5±0.097    |

\*p<0.0001; \*\*p<0.01; \*\*\*p<0.05 when compared with control

Paracetamol group significantly increased the serum level of SGOT (132.59%), SGPT (83.33%), Bilirubin (59.68%) and ALP (172.72%) as shown in table 2. The 't' value obtained exceeded the limit of significance. That is in all cases, p < 0.0001.

**Treatment Groups (Table 2)**

**Positive control (silymarin) Groups (G3):** There was significant increase in total Bilirubin (1.633 ± 0.206 mg/dl), accompanied by significant decrease in level of SGOT (63.17 ± 2.20 mg/dl) and also significant decrease in ALP (41.67 ± 1.80 IU/dl), SGPT (79.67 ± 1.69 IU/dl), as compared to the negative control.

**Ethyl acetate low dose Groups (G4):** There was significant increase in total Bilirubin (1.8 ± 0.110 mg/dl), accompanied by significant decrease in level of SGOT (89.83 ± 3.60 mg/dl) and also significant decrease in ALP (79.67 ± 1.43 IU/dl), SGPT (114.50 ± 2.05 IU/dl), as compared to the negative control.

**Ethyl acetate high dose (G5):** There was significant decrease in total Bilirubin (1.417 ± 0.130 mg/dl), accompanied by significant decrease in level of SGOT (84.33 ± 1.61 mg/dl) and also significant decrease in ALP (59.0 ± 1.88 IU/dl), SGPT (98.83 ± 1.45 IU/dl), as compared to the negative control.

**Ethanol low dose (G6):** There was significant increase in total Bilirubin (1.550 ± 0.123 mg/dl), accompanied by significant decrease in level of SGOT (80.50 ± 1.48 mg/dl) and also significant decrease in ALP (68.17 ± 1.89 IU/dl), SGPT (104.33 ± 1.87 IU/dl), as compared to the negative control.

**Ethanol high dose (G7):** There was significant decrease in total Bilirubin (1.4 ± 0.106 mg/dl), accompanied by significant decrease in level of SGOT (66.83 ± 1.08 mg/dl) and also significant decrease in ALP (48.0 ± 1.73 IU/dl), SGPT (86.83 ± 1.66 IU/dl), as compared to the negative control.

**Histopathological studies results**

Hepatocytes of the ethyl acetate extracts pretreated group, showed moderate inflammation with moderate cell death and their lobular architecture was normal. Hepatocytes of the ethanolic extracts pretreated group showed minimal inflammation with minimal cell death and their lobular architecture was normal.

**Table 2: Effects of ethyl acetate and ethanolic extracts of *Tecomaria capensis* plant on various biochemical parameters in rats with paracetamol induced hepatotoxicity.**

| Group            | SGOT        | SGPT         | ALP         | Total bilirubin | Direct      | Indirect       |
|------------------|-------------|--------------|-------------|-----------------|-------------|----------------|
| Control          | 53.67±1.91  | 72.0±1.39    | 33.0±1.53   | 0.95±0.099      | 0.667±0.067 | 0.7±0.106      |
| Negative control | 124.83±2.52 | 132.0±1.10   | 90.0±1.29   | 1.517±0.168     | 1.2±0.137   | 1.033±0.109    |
| Positive control | 63.17±2.20  | 79.67±1.69   | 41.67±1.80  | 1.633±0.206     | 0.583±0.114 | 0.817±0.095    |
| Ethylacetate     | 89.83±3.60* | 114.50±2.05* | 79.67±1.43* | 1.8±0.110**     | 0.833±0.088 | 1.100±0.106*** |
| Low dose         |             |              |             |                 |             |                |
| Ethylacetate     | 84.33±1.61* | 98.83±1.45*  | 59.0±1.88*  | 1.417±0.130***  | 0.883±0.135 | 0.7±0.106      |
| High dose        |             |              |             |                 |             |                |
| Ethanol          | 80.50±1.48* | 104.33±1.87* | 68.17±1.89* | 1.550±0.123**   | 0.717±0.135 | 1.033±0.150    |
| Low dose         |             |              |             |                 |             |                |
| Ethanol          | 66.83±1.08* | 86.83±1.66*  | 48.0±1.73*  | 1.4±0.106***    | 0.817±0.101 | 0.667±0.088    |
| High dose        |             |              |             |                 |             |                |


\*p<0.0001; \*\*p,0.01; \*\*\*p,0.05 when compared with control

**Histopathological results of ethyl acetate and ethanolic extracts of *Tecomaria capensis* plant**




**Fig. 1: Control**

Hepatocytes of the normal control group showed a normal lobular architecture of the liver.




**Fig. 2: Silymarin (STD)**

Hepatocytes of the silymarin treated group showed minimal inflammation with minimal cell death and their lobular architecture was normal.




**Fig. 3: Negative Control (CCl<sub>4</sub>)**

Hepatocytes of the CCl<sub>4</sub> treated group showed maximum inflammation with maximum cell death.




**Fig. 4: Negative Control (PARACETAMOL)**

Hepatocytes of the Paracetamol treated group showed maximum inflammation with maximum cell death




**Fig. 5: Ethyl Acetate High Dose**

**Hepatocytes of the ethyl acetate extract (high dose) pretreated group showed moderate inflammation with moderate cell death and their lobular architecture was normal.**




**Fig. 6: Ethyl Acetate Lowdose**

**Hepatocytes of the ethyl acetate extract (low dose) pretreated group showed moderate inflammation with moderate cell death and their lobular architecture was normal.**



**Fig. 7: Ethanol High Dose**

**Hepatocytes of the ethanolic extract (high dose) pretreated group showed minimal inflammation with minimal cell death and their lobular architecture was normal.**



**Fig. 8: Ethanol Low Dose**

**Hepatocytes of the ethanolic extract (low dose) pretreated group showed minimal inflammation with minimal cell death and their lobular architecture was normal.**

Liver damage induced by  $\text{CCl}_4$  is commonly used model for the screening of hepatoprotective drugs. The rise in serum levels of AST, ALT and cholesterol has been attributed to the damaged structural integrity of the liver, because they are cytoplasmic in location and released into circulation after cellular damages. When rats were treated with carbon tetrachloride, it induces hepatotoxicity by metabolic activation, therefore, it selectively causes toxicity in liver cells maintaining semi-normal metabolic function. Carbon tetrachloride is metabolically activated by the cytochrome P-450 dependent mixed oxidase in the endoplasmic reticulum to form trichloromethyl free radical ( $\text{CCl}_3$ ) which combined with cellular lipids and proteins in the presence of oxygen to induce lipid peroxidation. This results in changes of structures of the endoplasmic reticulum and other membrane, loss of metabolic enzyme activation, reduction of protein synthesis and loss of glucose-6-phosphatase activation, leading to liver injury. Treatment with ethyl acetate and ethanolic extract recovered the injured liver towards normal, so it is having hepatoprotective effect.

Paracetamol hepatotoxicity is caused by the reaction metabolite N-acetyl-p-benzo quinoneimine (NAPQI), which causes oxidative stress and glutathione depletion. It is a well-known antipyretic and analgesic agent, which produces hepatic necrosis at higher doses<sup>9</sup>.

Paracetamol toxicity is due to the formation of toxic metabolites, when a part of it is metabolized by cytochrome P-450. Introduction of cytochrome<sup>10</sup> or depletion of hepatic glutathione is a prerequisite for paracetamol induced hepatotoxicity<sup>11</sup>. Normally, AST and ALP are present in high concentration in liver. Due to hepatocyte necrosis or abnormal membrane permeability, these enzymes are released from the cells and their levels in the blood increases. ALT is a sensitive indicator of acute liver damage and elevation of this enzyme in non-hepatic diseases is unusual. ALT is more selectively a liver parenchyma enzyme than AST<sup>12</sup>. Assessment of liver function can be made by estimating the activities of serum ALT, AST, ALP and Bilirubin which are enzymes originally present in higher concentrations in cytoplasm. When there is heaptopathy, these enzymes leak into the blood stream in conformity with the extent of liver damage<sup>13</sup>. The elevated level of these entire marker enzymes observed in the group II, paracetamol treated rats in this present study corresponded to the extensive liver damage induced by toxin. The reduced concentrations of ALT, AST and ALP as a result of plant extract administration observed, during the present study, might probably be due in part to the presence of flavonoids. Liver protective herbal drugs contain a variety of chemical constituents like phenols, coumarins, lignans, essential oil, monoterpenes,

carotenoids, glycosides, flavonoids, organic acids, lipids, alkaloids and xanthenes<sup>14</sup>. Treatment with ethyl acetate and ethanolic extract recovered the injured liver towards normal, so it is having hepatoprotective effect. Comparing these two extracts of *Tecomaria capensis* ethanolic extract is showing more significant hepatoprotective action than ethyl acetate extract.

#### ACKNOWLEDGEMENT

The authors are thankful to the Management and Staff members of Vignan Pharmacy College, Vadlamudi, Guntur Dt, and A.P for their support.

#### REFERENCES

- Guntupalli M: Hepatoprotective effects of rubiadin, a major constituent of *Rubia cordifolia* Linn. J. Ethnopharmacol 2006; 103, 484-490
- Chatterjee TK. Medicinal plants with hepatoprotective properties. In: Herbal Options. 3rd Edn. Books and Allied (P) Ltd. Calcutta. 2000; 135
- South African National Biodiversity Institute's (SANBI) Integrated Biodiversity System, information URL:[http://www.plantzafrica.com/plantuv/tecomar\\_cap.htm](http://www.plantzafrica.com/plantuv/tecomar_cap.htm) (10 December 2010)
- Roberts, M., Indigenous Healing Plants. Halfway House. Southern Book Publishers, 1990
- Hutchings, A., Scott, A.H., Lewis, G., Cunningham, A.B., Zulu medical plants; An Inventory. University of Natal press, Pietermaritzburg, 1996
- P.Pillaya, V.J.Maharaj, P.J.Smith: Journal of ethanopharmacology 2008; 119, 438-454
- Gomase.V.,Rangari.D : Phyto chemical evaluation and hepatoprotective activity of fresh juice of young stem (Tender) bark of *Azadirachta indica* a. juss. International journal of pharmacy and pharmaceutical sciences, 2011; suppl 2: 55-59.
- Manokaran S, Jaswanth A, Sengottuvelu S, Nandhakumar J, Duraisamy R, Karthikeyan D2 and Mallegaswari R: Hepatoprotective Activity of *Aerva lanata* Linn. Against Paracetamol Induced Hepatotoxicity in Rats. *Research J. Pharm. and Tech.* 2008; 1-4: Oct.-Dec..
- Boyd EH and Berezky GM: Liver necrosis from paracetamol. *Br J Pharmacol* 1966; 26, 606-614
- Dahlin D et al. Nacetyl- p-benzoquinone imine: A cytochrome P-450- mediated oxidation product of acetaminophen. *Proc Natl Acad Sci.* 1984; 81, 1327-1331
- Moron MS, Depierre JW and Mannervik B: Levels of glutathione, glutathione reductase and glutathione-S-transferase activities in rat lung and liver. *Biochem Biophys Acta.* 1979; 582, 67-78.
- Shah M: Evaluation of the effect of aqueous extract from powders of root, stem, leaves and whole plant of *phyllanthus debilis* against CCL<sub>4</sub> induced rat liver dysfunction. *Indian Drugs.* 2002; 39, 333-337
- Nkosi CZ, Opoku AR and Terblanche SE: Effect of pumpkin seed (*Cucurbita pepo*) protein isolate on the activity levels of certain plasma enzymes in CCL<sub>4</sub>-induced liver injury in lowprotein fed rats. *Phy.the.Res.*2005; 19, 341-345
- Gupta AK and Misra N: Hepatoprotective Activity of Aqueous Ethanolic Extract of *Chamomile capitula* in Paracetamol Intoxicated Albino Rats. *American Journal of Pharmacology and Toxicology.* 2006; 1, 17-20