

ASSESSMENT OF RISK FACTORS AND MEDICATION ADHERENCE OF HYPERTENSION PATIENTS IN A TERTIARY CARE TEACHING HOSPITAL

V.HARISH KUMAR*, M. MOUNIKA, G. TEJA REDDY, S. MAHESH KUMAR

Department of clinical pharmacy, Smt Sarojini Ramulamma College of Pharmacy, Mahabubnagar-509001, Andhra Pradesh, India.
Email: harishvundhyala1989@gmail.com

Received: 28 Jan 2014, Revised and Accepted: 08 Mar 2014

ABSTRACT

Hypertension is a chronic condition, which leads to serious complications like cardiovascular diseases and stroke.

Objective: The main objective of this study was to assess the risk factors and med adherence for hypertensive patients in rural tertiary care teaching hospital by using risk assessment form and morisky medication adherence questionnaire form respectively.

Methods: These forms contains closed ended questionnaires. It is a prospective observational study conducted over a period of six months.

Results: In this study 278 subjects consisting of 150(54%) men and 128(46%) women had hypertension. In this study Age, gender, BMI, Family history, Smoking and Alcoholic were observed for risk factor assessment and morisky medication adherence scale is used to assess adherence of the patient towards medication. In hypertension patients, age is a major risk factor, in which more number of patients of age 51-60 years were 76(27.%) found. Commonly seen risk factors were age, gender, BMI, and family history. More no of patients were highly adherent to their medication i.e 55%.

Conclusion: By this study we concluded that more number of male hypertension patients were exposed to risk factors than female and more patients are highly adherent to their medication.

Keywords: Hypertension, Risk factors, Medication adherence.

INTRODUCTION

Risk factor refers to an attribute or characteristic or exposure of an individual whose presence or absence raises the probability of an adverse health outcome¹. Medication adherence² is defined as extent to which a patient's medication taking behaviour coincides with the intension of the health advice he or she has been given. Medication adherence is one of the important factors that determine therapeutic outcomes, especially in patients suffering from chronic illness. Currently non-communicable diseases, such as Diabetes, Asthma, COPD, Hypertension and Heart Disease are rapidly replacing infectious diseases. While mortality due to communicable diseases is decreasing, that for non-communicable diseases like chronic disease is rising at a very rapid pace. Chronic diseases are the leading cause of death in both males and females in all WHO regions. Approximately 72% of all chronic diseases occur in people aged 30 years and older. The causes of the main chronic diseases were risk factors. Some risk factors can be modified, other can't be modified. The modifiable factors include smoking, hypertension, elevated serum cholesterol, physical activity, obesity. The unmodifiable risk factors such as age, sex, race, family history and genetic factors³. Medication adherence is a important factor that determine therapeutic illness, various factors associated with the medication adherence are age, sex, annual income, education and social habits. Hypertension is classified into primary hypertension, which accounts for the majority of adulthood hypertension with no identifiable cause, although there are usually recognizable risk factors; and secondary hypertension, which accounts for the majority of childhood hypertension. Hypertension is the most common cardiovascular disorder affecting approximately 1 billion people globally and accounts for approximately 7.1 million deaths annually⁴.

MATERIALS AND METHODS

Study site

The Study was conducted in the General Medicine Department of SVS MEDICAL COLLEGE HOSPITAL, MAHABUBNAGAR which is a Tertiary Care Teaching Hospital which has 900 beds with Multi speciality departments.

Study design

The study was prospective observational and cross-sectional study.

Study period

This study was conducted over a period of six months starting from March 2013 to August 2013.

Study approval

This study was approved by the ethical committee constituted by SVS Medical College Hospital, Mahabubnagar.

Study Materials

Materials and Source of data

- Weighing machine
- Height measuring tape
- **Patient consent form-** An informed consent form was prepared in both English and Telugu version for the convenience of patient's understanding.
- **Patient data collection form-** It contains the socio demographic details of the patients like Age, Sex, BMI, Education, Occupation, Annual income, Smoking, Alcoholic, Family history details of HTN,DM,CKD.
- Medication adherence questionnaire
- Risk factors questionnaire

Statistical software

The Statistical software namely SAS 9.2, SPSS 15.0, Stata 10.1, MedCalc 9.0.1, Systat 12.0 and R environment ver.2.11.1 were used for the analysis of the data and Microsoft word and Excel have been used to generate graphs, tables etc.

Questionnaires

Morisky Medication Adherence Scale (MMAS) questionnaire

It is an 8 item Medication Adherence Scale of Morisky, prior permission was taken to use it before conducting the study. The MMAS is a generic self reported, medication taking behavior scale by which patient adherence to the medication is assessed. The

questionnaire was prepared in both Telugu and English version for convenience of patient's understanding and for providing information. It contains 8 questions with two options for first 7 questions and five options for 8 question.

STUDY PROCEDURE

This is a prospective observational study where patients eligible were enrolled into the study after obtaining the consent. The data collection form was prepared and used. This form mainly contains the demographic details of the patient. Questionnaires for risk factors and medication adherence were used to obtain the information about patient. All information relevant to the study was collected at the time of admission till the date of discharge and the data will be analysed using suitable method for statistical analysis. Medication adherence questionnaires consist of 8 questions by which adherence behaviour of patient can be assessed. Risk factors were assessed by knowing the Age, Sex, BMI, Occupation, Annual income, Family history, Comorbidities, Alcohol and Smoking history. The patients details were collected by using the case sheet or by directly asking the patients, the questionnaires were asked to be filled by the patients and the patients medication adherence, risk factors are assessed by the information

RESULTS AND DISCUSSION

A total number of 278 patients were enrolled for the study, among those male were 150(54%) and female were 128(46%) having hypertension as shown in Figure 1. Where as in the study conducted by Dogan N showed that female were more 374(31.3%) compared to that of male 119(14.10%)⁵.

According to the Table 1 among major six risk factors majority of the patients were having age risk factor By the collected data it is found that 51-60 years age group patients were more 76 (27.3%) as compare to remaining age groups. where as in a study conducted by Tazi MA et al. showed that 45-54 years age group patients were more 183(25.70%).

According to the Table 1 in BMI risk factor obese patients (BMI > 30 kg/m²) were 15(5.4%), over weight patients (BMI 23-30 kg/m²) were 126(45.3%), normal patients (BMI 18-23 kg/m²) were 126(45.3%) and under weight patients(BMI <18 kg/m²) were 11(4.0%). Among these normal BMI patients and over weight patients were more as compare to underweight and obese. The study conducted by Tazi MA et al. shown similar results that normal patients had 43.20% were more as compare to overweight 32.80% and obese patients 23.60%⁶.

According to the Table 1 in family history risk factor 181 patients (65.1%) were with the positive family history of HTN, DM, CKD i.e, 34.9%, 20.5%, 9.7% respectively and patients with no family history were 97(34.9%).

According to the Table 1 smokers were 80(28.8%) and 198(71.2%) were non-smokers. This study having similarities with the study conducted by Hazarika NC, in this study smokers were less 131(15.70%) as compared to the non-smokers 701(84.30%)⁷.

According to the Table 1 alcoholics were 106(38.1%) and 172(61.9%) were non-alcoholics this study having similarities with the study conducted by Hazarika NC, in this study alcoholics were less 328(36.94%) as compared to the non-alcoholics 560(63.60%)⁸.

Table 1: shows characteristics of the patients considered for the assessment of Risk factors


| Basic variables | No. of patients | % |
|-------------------------------|-----------------|--------------|
| Age in years | | |
| 25-30 | 24 | 8.6 |
| 31-40 | 34 | 12.2 |
| 41-50 | 57 | 20.5 |
| 51-60 | 76 | 27.3 |
| 61-70 | 59 | 21.2 |
| 71-80 | 21 | 7.6 |
| >80 | 7 | 2.5 |
| Gender | | |
| Male | 150 | 54.0 |
| Female | 128 | 46.0 |
| Smoking | | |
| Smokers | 80 | 28.8 |
| Non-Smokers | 198 | 71.2 |
| Alcoholic | | |
| Alcoholic | 106 | 38.1 |
| Non-Alcoholic | 172 | 61.9 |
| BMI (kg/m²) | | |
| <18.5 (under weight) | 11 | 4.0 |
| 18.5-23 (normal) | 126 | 45.3 |
| 23-30 (over weight) | 126 | 45.3 |
| >30 (obese) | 15 | 5.4 |
| Family history | | |
| Nil | 97 | 34.9 |
| Yes | 181 | 65.1 |
| • CKD | 27 | 9.7 |
| • DM | 57 | 20.5 |
| • HTN | 97 | 34.9 |
| Total | 278 | 100.0 |

Assessment of medication adherence is worked out based on the responses provide by the patients for the Morisky medication adherence questionnaires. The responses provided by the patients are analysed and reported in the Table 2 and the results are drawn from it and presented in Table 3. According to that among 278 patients 158(55.0%) were highly adherent, 68(24.5%) were medium adherent and 57(20.5%) were low adherent to their medications.

A Cross sectional study was conducted to analyse the medication adherence with respect to age, gender, education, income, and social habits.

According to Table 4 the highly adherent behaviour of patients with different age groups of 25-30, 31-40, 41-50, 51-60, 61-70, and 71-80 are 9.8%, 11.11%, 19.61%, 28.1%, 22.88%, and 8.5% respectively.

According to Table 4, 54.9% of the male patients and 45.1% of the female patients are highly adherent.



According to Table 4, the highly adherent behaviour of illiterates, primary educated, secondary educated and graduates are 38.56%, 9.8%, 32.68% and 18.95% respectively.

According to Table 4, the highly adherent behaviour of patients with the income groups <25000, 25000-50000, 50000-100000, and >100000 are 35.29%, 22.22%, 40.52%, and 1.96% respectively.

According to Table 4, the highly adherent behaviour of alcoholic patients and non- alcoholic patients are 37.25% and 62.75% respectively.

According to Table 4, the highly adherent behaviour of smokers and non- smokers are 20.26% and 79.74% respectively.

Table 2: Questionnaire analysis

| Questions | Response | |
|--|-------------|------------|
| | Yes (n=278) | No (n=278) |
| Do you sometimes forget to take your medicine? | 66(23.7%) | 212(76.3%) |
| Thinking over the past 2 weeks was there any days when you did not take your medicine? | 29(10.4%) | 249(89.6%) |
| Have you ever cut back or stopped taking your medicine without telling your doctor because you felt worse when you took? | 40(14.4%) | 238(85.6%) |
| When you travel or leave home do you sometimes forget to bring along your medicine? | 63(22.7%) | 215(77.3%) |
| Did you take all your medicine yesterday? | 248(89.2%) | 30(10.8%) |
| When you feel like your symptoms are under control do you sometimes stop taking your medicine? | 55(19.8%) | 223(80.2%) |
| Do you ever feel hassled about sticking to your treatment plan? | 74(26.6%) | 204(73.4%) |
| How often do you have difficulty remembering to take all your medicine? A. Never B.once in a while C.sometimes D.usually E.all the time (no= option 'A'; yes=option 'B-E') | 117(42.1%) | 161(57.9%) |

Table 3: shows the distribution of Morisky Medication Adherence Score

| Morisky Medication Adherence score | No. of patients | % |
|------------------------------------|-----------------|-------|
| 1-4 (low adherence) | 57 | 20.5 |
| 5-6 (medium adherence) | 68 | 24.5 |
| 7-8 (high adherence) | 153 | 55.0 |
| Total | 278 | 100.0 |

Table 4: shows the Correlation of Morisky medication adherence score according to age, gender, Education, income and social habits in HTN patients.

| Variables | Morisky medication adherence score | | | P value |
|----------------|------------------------------------|-------------------------------|------------------------------|---------|
| | Low Adherence (1-4) (n=57) | Medium Adherence (5-6) (n=68) | High Adherence (7-8) (n=153) | |
| Age in years | | | | 0.605 |
| • 25-30 | 6(10.53%) | 3(4.41%) | 15(9.8%) | |
| • 31-40 | 5(8.77%) | 12(17.65%) | 17(11.11%) | |
| • 41-50 | 14(24.56%) | 13(19.12%) | 30(19.61%) | |
| • 51-60 | 16(28.07%) | 17(25%) | 43(28.1%) | |
| • 61-70 | 8(14.04%) | 16(23.53%) | 35(22.88%) | |
| • 71-80 | 8(14.04%) | 7(10.29%) | 13(8.5%) | |
| Gender | | | | 0.708 |
| • Male | 28(49.12%) | 38(55.88%) | 84(54.9%) | |
| • Female | 29(50.88%) | 30(44.12%) | 69(45.1%) | |
| Education | | | | 0.073+ |
| • Illiterate | 28(49.12%) | 22(32.35%) | 59(38.56%) | |
| • Primary | 7(12.28%) | 13(19.12%) | 15(9.8%) | |
| • Secondary | 11(19.3%) | 14(20.59%) | 50(32.68%) | |
| • Graduate | 11(19.3%) | 19(27.94%) | 29(18.95%) | |
| Income | | | | 0.041* |
| • <25000 | 26(45.61%) | 19(27.94%) | 54(35.29%) | |
| • 25000-50000 | 16(28.07%) | 25(36.76%) | 34(22.22%) | |
| • 50000-100000 | 13(22.81%) | 20(29.41%) | 62(40.52%) | |
| • >100000 | 2(3.51%) | 4(5.88%) | 3(1.96%) | |
| Alcohol | | | | 0.836 |
| • Absent | 36(63.16%) | 40(58.82%) | 96(62.75%) | |
| • Present | 21(36.84%) | 28(41.18%) | 57(37.25%) | |
| Smoker | | | | 0.001** |
| • Absent | 30(52.63%) | 46(67.65%) | 122(79.74%) | |
| • Present | 27(47.37%) | 22(32.35%) | 31(20.26%) | |

CONCLUSION

The study results suggest that male patients, patients with the age group 51-60, over weight patients and patients with the positive family history were more exposed to risk factors when compared to other groups, more number of patients were highly adherent to their medications.

This study concludes that certain educational programs are needed to educate the people to emphasize the importance of medication adherence and modification to their life style to reduce the complications associated with hypertension.

ACKNOWLEDGMENT

The authors are thankful to Smt Sarojini Ramulamma College of Pharmacy, Mahabubnagar staff and management for their all time support. We are graceful to **Donald E Morisky** and **K.P.Suresh** for providing MMA Scale and statistical work respectively.

REFERENCES

1. Altobelli E, Petrocell R, Maccarone M, Altomare G, Argenziano G, Giannetti A et al. Risk factors of hypertension, diabetes and obesity in Italian psoriasis patients: a survey on socio-demographic characteristics, smoking habits and alcohol consumption. *Eur J Dermatol* 2009;19 (3):252-6.
2. url address: http://www.nhtsa.gov/people/injury/olddrive/druguse_older_driver/pages/FactorsAffecting.htm
3. Park K. Text book of Preventive and social medicine.16th ed: Jabalpur: Banarsidas Bhanot; 2000. P. 31, 272.
4. Wamala JF, Karyabakabo Z, Ndungutse D, Guwatudde D. Prevalence factors associated with Hypertension in Rukungiri District, Uganda-A Community-Based Study. *African Health Sciences* 2009;9(3): 153-60.
5. Dogan N, Toprak D, Demir S. Hypertension prevalence and risk factors among adult population in Afyonkarahisar region: a cross-sectional research. *Anadolu Kardiyol Derg* 2012;12:47-52.
6. Tazi MA, Tazi S, Khalil A, Lahmouz F, Arrach ML, Chaouki N. Risk factors for hypertension among the adult Moroccan population. *Eastern Mediterranean Health Journal* 2009;15(4):827-41.
7. Hazarika NC, Narain K, Biswas D, Kalita HC, Mahanta J. Hypertension in the native rural population of Assam. *THE NATIONAL MEDICAL JOURNAL OF INDIA* 2004; 17(6):300-04.
8. Hazarika NC, Biswas D, Mahanta J. Hypertension in the Elderly Population of Assam. *JAPI* 2003 june; 51:567-73.