

## THE PHARMACOLOGICAL IMPORTANCE OF ANETHUM GRAVEOLENS.A REVIEW

ALI ESMAIL AL-SNAFI

Department of Pharmacology, College of Medicine, Thi qar University, Nasiriyah, P O Box 42, Iraq. Cell: +9647801397994.

Email: aboahmad61@yahoo.com

Received: 24 Dec 2013 Revised and Accepted: 11 Jan 2014

### ABSTRACT

*Anethum graveolens* contained essential oils, fatty oil, proteins, carbohydrates, furanocoumarin, polyphenols, mineral and many other biologically active constituents. It is widely used traditionally. The pharmacological studies showed that *Anethum graveolens* induced antimicrobial, anti-inflammatory, analgesic, gastric mucosal protective and antisecretory effects, smooth muscle relaxant effect, hyperlipidaemic, increased progesterone concentration, and many other effects. This review will highlight the chemical constituent and pharmacological effects of *Anethum graveolens*.

**Keywords:** *Anethum graveolens*, Dill, chemistry, Pharmacology

### INTRODUCTION

*Anethum graveolens*, belong to the family Umbelliferae, is indigenous to southern Europe. It is an annual herb growing in the Mediterranean region, central and southern Asia. Now it is cultivated widely throughout the world [1-4]. It is used traditionally as a popular aromatic herb and spice that has a very long history of use going back to more than 5,000 years. It was used as a remedy for indigestion and flatulence and as milk secretion stimulant. Moreover, it is used as an anti-convulsion, anti-emetic, anti-cramp (in children), as a wound healer and to increase the appetite and strengthen the stomach [5-7].

*Anethum graveolens* contained essential oils, fatty oil, moisture (8.39%), proteins (15.68%), carbohydrates (36%), fiber (14.80%), ash (9.8%), furanocoumarin, polyphenols and mineral [6-13].

Previous studies showed that *Anethum graveolens* induced antimicrobial, anti-inflammatory, analgesic, gastric mucosal protective and antisecretory effects, smooth muscle relaxant effect, hyperlipidaemic, increased progesterone concentration, and many other effects. The aim of the present review is to highlight the chemical constituent and pharmacological effects of *Anethum graveolens*.

**Synonyms:** *Pastinaca anethum* Spreng, *Peucedanum graveolens* Benth. and *Selinum anethum* Roth.

**Common names:** Unani: Soyaa, Siddha: Sadakuppai, Sanskrit: Sthathpushpi, Hindi: Sowa, Punjabi: Soya, English: Dill and Anet, Arabic: Shibth and Haba helwa [7].

**Parts used medicinally:** Fruit, leaf, and essential oil

**Physicochemical properties** (4, 8-9):

Foreign organic matter: not more than 2.0%, total ash: not more than 11.0%, acid-insoluble ash: not more than 1.5%, water-soluble extractive: not less than 15.0%, and alcohol-soluble extractive: not less than 4.0%.

**Chemical constituents**

*Anethum graveolens* contained essential oils, fatty oil, moisture (8.39%), proteins (15.68%), carbohydrates (36%), fiber (14.80%), ash (9.8%) and mineral elements such as calcium, potassium, magnesium, phosphorous, sodium, vitamin A and niacin. Fruits of *Anethum graveolens* contain 1 - 4% essential oil comprising of major compounds: carvone (30 - 60%), limonene (33%),  $\alpha$ -phellandrene (20.61%), including pinene, diterpene, dihydrocarvone, cineole, myrcene, paramyrcene, dillapiol, isomyristicin, myristicin, myristin,

apiol and dillapiol. *Anethum graveolens* essential oil also contained furanocoumarin, 5-(4'-hydroxy-3"-methyl-2"- butenyloxy)-6, 7-furocoumarin, oxypeucedanin, oxypeucedanin hydrate and falcariindiol [6, 10-12].

The total phenol and total flavonoid contents of *Anethum graveolens* L. extract were 105.2 mg of gallic acid equivalents/g of the dried extract and 58.2 mg of catechin equivalents/g of the dried extract, respectively [13].

**Pharmacological effects**

**Antimicrobial effects**

The essential oil and different extracts of *Anethum graveolens* seeds exerted antimicrobial activity against wide range of microorganisms. The essential oils and acetone extracts shown antimicrobial activity against *Staphylococcus aureus*, *Bacillus cereus*, *Enterococcus faecalis*, *Listeria monocytogenes*, *Escherichia coli*, *Yersinia enterocolitica*, *Salmonella choleraesuis*, *S. typhimurium*, *Shigella flexneri*, *Salmonella typhi*, *Pseudomonas aeruginosa*, and *Mycobacterium*. *Anethum graveolens* seed extracts have also been reported to possess anti-ulcer activity, and have shown moderate activity against *Helicobacter pylori*. Aqueous and organic extracts of seeds have exhibited potent antibacterial activity. The essential oils also active against three fungi (a yeast, *Candida albicans* and two molds, *Penicillium islandicum* and *Aspergillus flavus*). D-limonene and D-carvone, have exhibited strong antifungal activity against *Aspergillus niger*, *Saccharomyces cerevisiae* and *Candida albicans*. Many authors mentioned that the antimicrobial activities could be attributed to furanocoumarin in *Anethum graveolens* [11, 14-22].

**Anti-inflammatory and analgesic effects**

The hydro alcoholic extract of the *Anethum graveolens* seed caused significant decrease in the inflammation and pain in rats [23].

*Anethum graveolens* oil and diclofenac-gel showed a significant ( $p < 0.001$ ) decrease in the paw volume in rats compared to the blank group. *Anethum graveolens* oil showed even more decrease in the paw volume compared to the diclofenac [24].

A single topical application of an ethanol extract of the fruits to the inner and outer surface of the ear of mice inhibited ear inflammation induced by 12-*O*- e-tradecanoylphorbol- 13 acetate by 60% [25].

A 10% aqueous extract of the fruits and 5% aqueous solution of the essential oil had analgesic effects in mice pain induced by hot plate and acetic acid writhing models. The effect of the fruits (1.0 g/kg body weight) was comparable to 200 mg / Kg body weight of acetyl salicylic acid [26].

### Effects on gastrointestinal system:

*A. graveolens* seed extracts possessed significant mucosal protective and antisecretory effects in the gastric mucosa lesions induced in mice by oral administration of HCl (1 N) and absolute ethanol. The acidity and total acid content were reduced by the orally or intraperitoneally administration of the extracts [27]. *Anethum graveolens* seed extracts exerted moderate activity against *Helicobacter pylori* [28]. The essential oil of *Anethum graveolens* reduced contractions of rabbit intestine [29]. Ethanol extract inhibited acetylcholine and histamine induced contractions of guinea-pig ileum [30]. Dill seeds have been used as household remedy to relief digestive problems such as stomachache, indigestion and flatulence. Dill water is believed to have a soothing effect and is given to babies to treat gripe, relieve hiccups and colic [7]. The essential oil was a mild carminative and reduced foaming *in vitro* [31].

### Hyperlipidaemic effects

The crude extract of *Anethum graveolens* L showed anti-hypercholesterolaemic and anti-hyperlipidaemic activities. The crude extracts of *A. graveolens* L. besides having strong anti-hyperlipidaemic effects, it improved the biological antioxidant status by reducing lipid peroxidation in liver and modulating the activities of antioxidant enzymes in rats fed with high fat [32-34]. Treatment of hyperlipidaemic rats with defatted ethanolic *Anethum graveolens* L. extract (single daily dose of 1 ml, equivalent to 500 mg of the plant powder) and high-fat diet for up to 10 and/or 30 days reversed the serum lipid levels compared to rats which were fed only high-fat diet. In addition, it induced significant increase in HMG-CoA/mevalonate ratio as compared to rats which were fed high-fat diet after treatment with defatted ethanolic *Anethum graveolens* L. extract for 30 days [13]. Dill powder and its essential oils also exerted hypolipidaemic activity in rats [34]. Intravenous administration of 12.5 mg/kg body weight of 70% dried ethanol extract of the fruits dissolved in normal saline or 4.0 µl/kg body weight of the essential oil induced diuresis and enhance sodium and calcium excretion in dogs [35]. Intravenous administration of 5-10 mg/kg body weight of 5% seed oil in saline to cats caused hypotension and increased respiration volume [2-4].

### Effects on reproductive system

The effects of *Anethum graveolens* L. (dill) extracts on female reproductive system were studied in female rats. The experimental groups were fed 0.045 g/kg and 0.45 g/kg of aqueous extract and 0.5 g/kg and 5 g/kg of ethanol extract for 10 days. Treatment with high dose of the extract resulted in a significant increase in duration of the estrous cycle and diestrus phase. Smooth endoplasmic reticulum (SER), rough endoplasmic reticulum (RER) and mitochondria were increased in granulosa lutein cells in high dose groups. There were no significant statistical differences in amount of serum estradiole between experimental, control and sham groups but the serum progesterone concentration increased significantly in high dose treatment group compared with control and sham groups [36].

Dill seed possessed contractive effects on myometer, enhanced releasing of oxytocin which is an effective hormone in uterus contractions. A dose of 6-7 gm of dill seed extract after delivery decreases postpartum hemorrhage due to its contractive characteristic. Limonene and anethole showed contractive effect on uterine myometrium [37-40].

Zagamil *et al.* carried out a clinical study to evaluate the effect of Dill seed on uterus contractions in active phase of labor. 40 women used Dill seed infusion (one tablespoon of whole dill seed seeped in a half or whole cup boiling water for 3-4 min before going to the hospital at the beginning of uterus contractions), and 60 women used nothing in the control group. Interpretable electronic fetal monitoring was obtained for half an hour at the beginning of the active phase. The Fall: Rise ratio was calculated by measuring the duration of time for a contraction to return to its baseline from its peak (fall) divided to the duration of its rise time to its peak (rise). The number of contractions in the treated group was

significantly more than the control group. The ratio of contraction's fall time to its rise time in the treated group was shorter than the control group. The study showed that dill seed shortens duration of the first stage of labor [41].

### Contraindication and adverse effects

It was safe, but rarely and in sporadic cases, it caused allergic reactions, oral pruritus, tongue and throat swelling, urticaria, vomiting and diarrhea. It was not recommended during pregnancy. The maximum non-fatal doses of aqueous and ethanolic extracts in mice were 0.45 g/kg and 5 g/kg (ip), respectively. LD50 values of the aqueous and ethanolic extracts were 3.04 g/kg, (ip) and 6.98 g/kg, (ip) respectively [27, 42-44].

### Dosage

The average daily dose of the seeds is 3 g; essential oil is 0.1-0.3g [5, 44-45].

### CONCLUSION

*Anethum graveolens* is a plant with wide range of chemical constituents which exerted many pharmacological effects. There is a great promise for development of novel drugs from *Anethum graveolens* to treat human diseases as a result of its effectiveness and safety.

### REFERENCES

1. Wren R C. Potter's new cyclopedia of botanical drugs and preparations. Saffron Walden, CW Daniel 1988.
2. Leung A Y, Foster S. Encyclopedia of common natural ingredients used in food, drugs and cosmetics. New York, John Wiley and Sons 1996.
3. Yazdanparast R, Bahramikia S. Evaluation of the effect of *Anethum graveolens* L. crude extracts on serum lipids and lipoproteins profiles in hypercholesterolaemic rats. DARU 2008; 16(2):88-94.
4. African pharmacopoeia. Vol.1. Lagos, Organization of African Unity, Scientific Technical and Research Commission 1985.
5. Zargari A. Medicinal Plants. 6<sup>th</sup> ed. Vol. II, Tehran University Press, Tehran 1996; pp 531-528.
6. Ishikawa T M, Kudo M, Kitajima J. Water-soluble constituents of dill. Chem Pharm Bull 2002; 55:501-507.
7. Kaur G J, Arora D S. Bioactive potential of *Anethum graveolens*, *Foeniculum vulgare* and *Trachyspermum ammi* belonging to the family Umbelliferae - Current status. Journal of Medicinal Plants Research 2010; 4(2): 87-94.
8. The Ayurvedic pharmacopoeia of India. Part I. Vol. II. New Delhi, Ministry of Health and Family Welfare, Department of Indian System of Medicine and Homeopathy 1999.
9. WHO monographs on selected medicinal plants, Vol.3. WHO Library Cataloguing in Publication Data. WHO 2007 Pp. 34-41.
10. Khafagy S M, Mnajed H K. Phytochemical investigation of the fruit of Egyptian *Anethum graveolens*. I. Examination of the volatile oil and isolation of dillapiole. Acta Pharmaceutica Suecica 1968; 5:155-162.
11. Stavri M, Gibbons S. The antimycobacterial constituents of Dill (*Anethum graveolens*). Phytother Res 2005; 19: 938-941.
12. Radulescu V, Popescu M L, Ilies D C. Chemical composition of the volatile oil from different plant parts of *Anethum graveolens* L. (Umbelliferae). Farmacia 2010; 58:594-600.
13. Yazdanparast R, Bahramikia S. Evaluation of the effect of *Anethum graveolens* L. crude extracts on serum lipids and lipoproteins profiles in hypercholesterolaemic rats. DARU 2008; 16(2):88-94.
14. Delaquis P J, Stanich K, Girard B. Antimicrobial activity of individual and mixed fractions of dill, cilantro, coriander and eucalyptus essential oils. Int J Food Microbiol 2002; 74: 101-109.
15. Singh G, Kapoor I P S, Pandey S K. Studies on essential oils: part 10: Antibacterial activity of volatile oils of some spices. Phytother Res 2001; 16: 680-682.
16. Lopez P, Sanchez C, Batlle R, Nerin C. Solid- and vapor-phase antimicrobial activities of six essential oils: susceptibility of

- selected food borne bacterial and fungal strains. J Agric Food Chem 2005; 53(17): 6939-6946.
17. Mohsenzadeh M. Evaluation of antibacterial activity of selected Iranian essential oils against *Staphylococcus aureus* and *Escherichia coli* in nutrient broth medium. Pak J Biol Sci 2007; 10: 3693-3697.
  18. Kaur G J and Arora D S. *In vitro* antibacterial activity of three plants belonging to the family Umbelliferae. Int J Antimicrob Agents 2008; 31:393-395.
  19. Kaur G J, Arora D S. Antibacterial and phytochemical screening of *Anethum graveolens*, *Foeniculum vulgare* and *Trachyspermum ammi*. BMC Complement. Altern. Med.2009, 9: 30.
  20. Arora D S, Kaur G J. Antibacterial activity of some Indian medicinal plants. J Nat Med 2007; 61:313-317.
  21. Jirovetz L, Buchbauer G, Stoyanova AS. Composition, quality control and antimicrobial activity of the essential oil of long time stored dill (*Anethum graveolens* L.) seeds from Bulgaria. J Agric Food Chem 2003; 18: 3854-3857.
  22. Rifat-uz-Zaman M S, Akhtar M S, Khan M S. *In vitro* antibacterial screening of *Anethum graveolens* L. Fruit, *Cichorium intybus* L. leaf, *Plantago ovata* L. seed husk and *Polygonum viviparum* L. root extracts against *Helicobacter pylori*. Int J Pharmacol 2006; 2:674-677.
  23. Valady A, Nasri S, Abbasi N. Anti-inflammatory and analgesic effects of hydroalcoholic extract from the seed of *Anethum graveolens* L. J Med Plants 2010; 9: 130-124.
  24. Naseri M, Mojab F, Khodadoost M. The study of anti-inflammatory activity of oil-based dill (*Anethum graveolens* L.) extract used topically in formalin-induced inflammation male rat paw. Iranian Journal of Pharmaceutical Research 2012; 11 (4): 1169-1174.
  25. Okuyama T *et al.* Studies on cancer bio-chemoprevention of natural resources. X. Inhibitory effect of spices on TPA-enhanced 3H-choline incorporation in phospholipids of C3H10T1/2 cells and TPA-induced mouse ear edema. Zhonghua Yaoxue Zazhi 1995; 47:421-430.
  26. Racz-Kotilla E, Rotaru G, Racz G. Anti-nociceptive effect of dill (*Anethum graveolens* L.). Fitoterapia 1995; 2:80-81.
  27. Hosseinzadeh H, Karimi K R, Ameri M. Effects of *Anethum graveolens* L. seed extracts on experimental gastric irritation models in mice. Pharmacol 2002; 2: 21.
  28. Rifat-uz-Zaman, Akhtar M S, Khan M S. *In vitro* antibacterial screening of *Anethum graveolens* L. Fruit, *Cichorium intybus* L. leaf, *Plantago ovata* L. seed husk and *Polygonum viviparum* L. root extracts against *Helicobacter pylori*. Int J Pharmacol 2006; 2:674-677.
  29. Shipochliev T. Pharmacological investigation into several essential oils. I. Effect on the smooth musculature. Veterinarno Meditsinski Nauki 1968; 5:63-69.
  30. Dhar M L, Dhar M M, Dhawan B N, Mehrotra A C R. Screening of Indian plants for biological activity: part I. Indian Journal of Experimental Biology 1968; 6:232-247.
  31. Harries N, James K C, Pugh W K. Antifoaming and carminative actions of volatile oils. Journal of Clinical Pharmacology 1978; 2:171-177.
  32. Yazdanparast R, Bahramikia S. Improvement of liver antioxidant status in hyper- cholesterolaemic rats treated with *A. graveolens* extracts. Pharmacologyonline 2007; 3:88-94.
  33. Yazdanparast R, Alavi M. Antihyperlipidaemic and antihypercholesterolaemic effects of *Anethum graveolens* leaves after the removal of furocoumarins. Cytobios 2001; 105:185-191.
  34. Hajhashemi V, Abbasi N. Hypolipidemic activity of *Anethum graveolens* in rats. Phytother Res 2008; 22: 372-375.
  35. Mahran G H, Kadry H A, Isaac Z G, Thabet C K, Al-Azizi M M, El-Olemy M M. Investigation of diuretic drug plants. 1- Phytochemical screening and pharmacological evaluation of *Anethum graveolens* L., *Apium graveolens* L., *Daucus carota* L. and *Eruca sativa* Mill. Phytotherapy Research 1991; 5:169-172.
  36. Monsefi M, Ghasemi M and Bahaoddini A. The effects of *Anethum graveolens* L. on female reproductive system of rats. DARU 2006; 14 (3):131-135.
  37. Bertram G Z, Emertius P H. Basic and Clinical Pharmacology, 8<sup>th</sup> ed, New York Mc Graw Hill 2001.
  38. Mahdavian M, Golmakani N, Mansoori H, Afzalaghaee M. An investigation of effectiveness of oral dill extracts on postpartum hemorrhage. The Iranian Journal of Obstetrics, Gynecology and Infertility 2001; 4 (8-7):26-19.
  39. Gharibn Aseri M K, Mard S A, Farboud Y. Effect of *Anethum graveolens* fruit extract on rat uterus contractions. Iranian J Basic Med Sci 2005; 8(4 (28)): 263-270.
  40. Committee for Veterinary Medicinal Products, EMEA/MRL "Juniperi Fructus is the dried berry like cones of *Junipers communis*", 1999.
  41. Zagami S E, Golmakani N, Kabirian M, Shakeri M T. Effect of Dill (*Anethum graveolens* Linn.) seed on uterus contractions pattern in active phase of labor. Indian Journal of Traditional Knowledge 2012; 11 (4): 602-606.
  42. Chui A M, Zacharisen M C. Anaphylaxis to dill. Annals of Allergy, Asthma and Immunology 2000; 84:559-560.
  43. Nath D, Sethi N, Singh R K, Jain A K. Commonly used Indian abortifacient plants with special reference to their teratologic effect in rats. Journal of Ethnopharmacology 1992; 36:147-154.
  44. PDR for herbal medicines, Medical Economic Co. Montvale, New Jersey, 1998, p647.
  45. Blumenthal M Blumenthal M, Busse WR, Goldberg A (eds). The complete German Commission E monographs. Austin, TX, American Botanical Council, 1998.