

Research Article

QUANTIFICATION OF PRO-VITAMIN A ACTIVITIES AND CONTENT IN 22 SELECTED 'ULAM' SPECIES OR MALAYSIAN TRADITIONAL VEGETABLES

¹FATIMAH AZZHARA MOHD ZAIFUDDIN, ¹NORAZIAN MOHD HASSAN, ^{2*}RASHIDI OTHMAN

¹Department of Pharmaceutical Chemistry, Kulliyah of Pharmacy, International, Islamic University Malaysia, Kuantan, Malaysia,

²International Institute for Halal Research and Training (INHART), Herbarium Unit, Department of Landscape Architecture, Kulliyah of Architecture and Environmental Design, International Islamic University Malaysia, Kuala Lumpur, Malaysia.

Email: rashidi@ium.edu.my, adira_27@yahoo.com

Received: 05 Feb 2014 Revised and Accepted: 27 Mar 2014

ABSTRACT

Recently there is a considerable interest in combating vitamin A deficiency (VAD) by manipulating carotenoid content and composition in plants through genetic manipulation to meet the daily requirements for vitamin A. In Malaysia and most developing countries many people survive largely on plant-based diets or monotonous consumption such as cereals and legumes that are poor sources of vitamin A. *Ulam* or traditional vegetables consumed by Malaysian comprise more than 120 species representing various plant leaves, fruits, seeds, tubers and roots are valuable sources of nutrients and pro-vitamin A. These pro-vitamin A compounds can profoundly protect our well-being or risk of disease throughout our lives. Therefore, identifying nutritional quality of food crops and its ingredients for VAD is one of the urgent health issues and research worldwide. This study established that traditional vegetables or *ulam* differ greatly with respect to types and concentrations of pro-vitamin A. A total of 22 *ulam* species were evaluated for quantitative and qualitative pro-vitamin A composition through HPLC analysis. Three main pro-vitamin A carotenoids were identified namely β -cryptoxanthin, α -carotene and β -carotene. These pro-vitamin A groups content and ratio varies between *ulam* species but were detected higher in *cekur manis* ($68.44 \pm 0.23 \mu\text{g/g DW}$), *pegaga* ($2142.71 \pm 11.70 \mu\text{g/g DW}$) and *daun selom* ($3085.02 \pm 5.87 \mu\text{g/g DW}$). Total retinol equivalent activity presented by these pro-vitamin A groups in every *ulam* species demonstrated that *beluntas* has the highest activity ($656.59 \mu\text{g/g DW}$). The information gathered in this study embarked that almost all of the *ulam* species are good and potent sources of pro-vitamin A.

Keywords: pro-vitamin A, Carotenoid, β -cryptoxanthin, α -carotene, β -carotene and *ulam*.

INTRODUCTION

Vitamin A deficiency (VAD) has been pronounced as an urgent public health issue which is affecting almost all nations worldwide especially in low-income and developing countries[1]. This deficiency had caused major mortality among children aged below five years old and symptoms ranging from night blindness, xerophthalmia to those of keratomalacia which lead to the total blindness. Without adequate vitamin A supplementations, the children are not only having risks to be blind but also vulnerable to various kinds of diseases[2]. Therefore, improved dietary vitamin A status would be important for the population survival especially among the children thus eliminating VAD as a continuous public health concern[3]. Dietary vitamin A may present in two forms; pre-formed vitamin A from animal-based foods and pro-vitamin A that usually originating from plant-based foods. Pro-vitamin A that comes from photosynthetic plants and bacteria is collectively known as carotenoids[4,5]. They are lipid soluble pigments of yellow-orange-red colour which primarily act as antioxidant in photosynthetic tissues. Even though they are yellow-to-red coloured pigments, they are found concentrated in green leafy vegetables[6-10]. There are more than 700 carotenoids have been characterised where 40 of them were found in human diet such as α -carotene, β -carotene, lutein and lycopene[11]. Consumption of carotenoids in human diet has been reported to be associated with better health status and reduced fatality due to chronic diseases[11-16].

There are more than 120 species of traditional vegetables or *ulam* in Malaysia. They are important to the local community for their own unique taste in various dishes as well as their health benefits[17]. They are reported to be nutritious as they are rich in carbohydrates, proteins, minerals and vitamins. Some of the vegetables are claimed to have medicinal properties in such way they are beneficial in lowering the incidence of cancer and age-related diseases. They are also popular for their anti-ageing properties[18,19]. Despite of their popularity, scientific reports on their medicinal properties especially regarding the ones related to pro-vitamin A carotenoids' activities are still lacking. Therefore, this research is primarily aimed to explore alternative sources for pro-vitamin A carotenoids from various *ulam* species.

MATERIALS AND METHODS

Sample preparation

Edible parts of all samples were freeze-dried (EYELA FDU-1100, Japan) for 72 hours, after which the samples were ground into fine powder and kept at -20°C until further analysis.

Extraction of carotenoids

0.1 g of each powdered sample was rehydrated with adequate distilled water and extracted with a mixture of acetone and methanol (7:3) until the solution became colourless. Equal amount of hexane and distilled water were added to the combined solution to extract out the carotenoids. After separation of layers was achieved, the upper hexanal layer was collected and dried down under a gentle stream of oxygen-free nitrogen[21].

Saponification

Samples were saponified with a mixture of acetonitrile and water (9:1) and methanolic potassium hydroxide solution (10% w/v). Base carotenoids were then extracted out with hexane and 0.1% butylated hydroxyl toluene (BHT), followed by addition of 10% NaCl until phase separation was achieved. After being washed, the extracts were dried under a gentle stream of oxygen-free nitrogen. All of the extraction and saponification procedures were detailed in our previous work[20]. Finally, the dried extracts were re-suspended in ethyl acetate prior to HPLC analysis.

HPLC analysis

The HPLC analysis of saponified extracts was performed on an Agilent model 1200 series as detailed by Othman[21]. The column used was a ZORBAX SB-C₁₈ end capped $5\mu\text{m}$, $4.6 \times 250 \text{ mm}$ reverse phase column (Agilent Technologies, USA). The eluents used were (A) acetonitrile:water (9:1 v/v) and (B) ethyl acetate. The column separation was allowed via a series of gradient at a flow rate of 1.0 mL min^{-1} . The column then was allowed to re-equilibrate in 100% A for 10 min prior to the next injection.

Throughout the analysis, temperature of the column was maintained at 20°C while the injection volume was 10µL each. Carotenoid standards for β-carotene and β-cryptoxanthin were obtained commercially from Sigma-Aldrich and α-carotene standard was isolated from orange-fleshed carrots as described by Rodriguez-Amaya and Kimura[22]. Each of the individual carotenoids was detected at the wavelengths of 454 nm for β-cryptoxanthin and β-carotene (maximum absorption for both carotenoids in the mobile phase) and 447 nm for α-carotene. Compounds from the extracts were identified by co-chromatography with standards and by elucidation of their spectral characteristics using a photo-diode array detector.

The total and individual carotenoid concentrations were expressed in terms of microgram per gram dry weight of freeze-dried matter (µg/g DW).

Calculation of vitamin A activity

The vitamin A activity of each of the detectable pro-vitamin A carotenoids was converted into retinol equivalent units as proposed

by NRC[23], where 6 µg of β-carotene or 12 µg of α-carotene and β-cryptoxanthin is equal to 1 µg retinol equivalent (RE).

RESULTS AND DISCUSSION

A group of 22 species of commonly consumed Malaysian *ulam* (Table 1) had been selected for identification and quantification of their pro-vitamin A carotenoid content. As to date, there is no previous report on the carotenoid content for the following five *ulam* species (Table 1): *Daun Kesum*, *Hempedu Bumi*, *JarumTujuh Bilah*, *Kemangi* and *Mas Cotek*. Therefore, the results included in this report would provide new information for the said species with regard to their carotenoid accumulation.

Throughout this study, all of the experimental procedures were performed under dim light environment since the carotenoids are light-sensitive which otherwise would trigger to be photo-isomerised and photo-oxidised. In such case, analysis may come out with false positive results in the presence of carotenoid artifacts. Wherever needed; the extracts were completely dried under oxygen-free nitrogen gas prior to storage as to prevent individual carotenoid losses.

Table 1: Selected traditional vegetables (*ulam*) used in this study

Local name	Botanical name	Carotenoids detected in previous studies	References
Beka	<i>Oroxylum indicum</i>	β-carotene, lutein, violaxanthin	20
Beluntas	<i>Pluchea indica</i>	β-carotene, lutein, violaxanthin	20, 24
Cekur Manis	<i>Sauropus androgynus</i>	β-carotene, zeaxanthin, lutein, violaxanthin, neoxanthin	7, 9, 20, 24, 31
Daun Bawang	<i>Allium cepa</i>	β-carotene	10, 31
Daun Kari	<i>Murraya koenigii</i>	β-carotene, zeaxanthin, lutein, violaxanthin, neoxanthin	7, 10, 29, 31, 33
Daun Kesum	<i>Polygonum minus</i>	N/A	N/A
Daun Kucai	<i>Allium tuberosum</i>	β-carotene	7
Daun Selom	<i>Oenanthe javanica</i>	β-carotene, lutein	20
Daun Sup	<i>Apium graveolens</i>	β-carotene	25, 31
Gajus	<i>Anacardium occidentale</i>	β-carotene, lutein	7, 20, 24
Hempedu Bumi	<i>Andrographis paniculata</i>	N/A	N/A
JarumTujuh Bilah	<i>Pereskia sacharosa</i>	N/A	N/A
Kaduk	<i>Piper sarmentosum</i>	β-carotene, lutein	20, 25
Kemangi	<i>Ocimum americanum</i>	N/A	N/A
Mas Cotek	<i>Ficus deltoidea</i>	N/A	N/A
Salad	<i>Lactuca sativa</i>	β-carotene, zeaxanthin, lutein, violaxanthin, neoxanthin	10, 27, 32, 29, 30
Sawi	<i>Brassica chinensis</i>	β-carotene, zeaxanthin, lutein	7, 25, 26, 31
Selasih	<i>Ocimum basilicum</i>	β-carotene	25
Sirih	<i>Piper betle</i>	β-carotene, zeaxanthin, lutein	10, 29
Pegaga	<i>Centella asiatica</i>	β-carotene, lutein, violaxanthin, neoxanthin	20, 24, 25
TenggekBurung	<i>Euodia redlevi</i>	β-carotene, lutein	20
Ulam Raja	<i>Cosmos caudatus</i>	β-carotene, zeaxanthin, lutein	20, 24

N/A- Not available

Table 2: Pro-vitamin A activities and content in 22 selected traditional vegetables (*ulam*).

Local Name	BCR (µg per g DW)	AC	BC	Total (µg per g DW)
Salad	nd	nd	542.22±1.63	542.22±1.63
Kemangi	nd	nd	783.43±0.05	783.43±0.05
Mas Cotek	nd	379.35±0.06	874.62±0.04	1253.97±0.13
Tenggek Burung	nd	nd	1259.79±2.82	1259.79±2.82
Kaduk	nd	nd	1268.56±0.10	1268.56±0.10
Beka	nd	384.1 ±2.87	1123.96±3.42	1508.07±5.98
Daun Kari	nd	nd	1690.11±0.09	1690.11±0.09
Sawi	45.11±0.04	nd	1683.62±25.90	1728.73±25.94
Gajus	nd	nd	1738.12±0.09	1738.12±0.09
Hempedu Bumi	nd	nd	1766.17±2.45	1766.17±2.45
JarumTujuh Bilah	nd	nd	1863.58±15.55	1863.58±15.55
Daun Bawang	nd	nd	2066.85±2.45	2066.85±2.45
Selasih	nd	530.25±0.04	1950.47±24.11	2480.72±24.10
Daun Kucai	62.45±0.29	752.64±8.45	1856.99±0.09	2672.08±8.69
Sirih	67.64±0.34	nd	2739.45±9.88	2807.09±10.14
Daun Sup	58.99±0.03	nd	2757.76±8.90	2816.75±8.91
Ulam Raja	nd	1561.99±16.50	1430.11±5.17	2992.10±11.64
Daun Selom	nd	nd	3085.02±5.87	3085.02±5.87
Daun Kesum	nd	710.12±7.87	2532.96±25.38	3243.08±32.72
Pegaga	nd	2142.71±11.70	2387.64±6.20	4223.15±17.63
Cekur Manis	68.44±0.23	1357.80±0.14	2841.60±0.12	4267.84±0.45
Beluntas	nd	1915.74±15.58	2981.69±0.10	4897.43±15.51

nd- not detected, BCR- β-cryptoxanthin, AC- α-carotene, BC- β-carotene

Data represented as mean±standard deviation, n=5, significantly different at $p<0.0001$.

Total carotenoids are the sum of each pro-vitamin A carotenoid detected.

Under the experimental condition employed, it was observed that there was certain degree of variation in the level of total and individual carotenoid content among the selected species, ranging from one to nine fold (Table 2).

Among the species studied, HPLC analysis demonstrated that *Beluntas* accumulated substantially the highest total carotenoid (4897.43±15.51 µg/g DW) followed by *Cekur Manis* and *Pegaga* with 4267.84±0.45 µg/g DW and 4223.15±17.63 µg/g DW respectively. The total carotenoid content in all of the analysed samples was comparable with reports from previous studies. Interestingly, the total pro-vitamin A carotenoid concentrations for eight of the samples (*Tenggek Burung*, *Kaduk*, *Gajus*, *Ulam Raja*, *Daun Selom*, *Pegaga*, *Cekur Manis* and *Beluntas*) were found comparatively greater than the ones analysed in previous work elsewhere[20]. On top of that, the total pro-vitamin A carotenoid content for *Salad* obtained here was quantified lower than the data reported by Burns *et al.*[27] yet higher than values obtained by Mamatha[32].

The HPLC analysis also established that in determination of three types of pro-vitamin A carotenoids; β-cryptoxanthin which is a rare carotenoid, had been successfully quantified in *Sawi*, *Daun Sup*, *Daun Kucai*, *Sirih* and *Cekur Manis*. In fact, up to present date there is none of the previous analysis on the selected vegetables and herbs species reported on their β-cryptoxanthin content. Therefore, the results from this study established new profiling for this particular carotenoid.

α-carotene and β-carotene are isomeric compounds which differ in the location of double bond in one of their rings (Figure 1). Among the 22 *ulam* species analysed, β-carotene had been detected as their major pro-vitamin A compound but α-carotene was present in only nine species. This phenomenon is probably due to their function and

the regulation of their biosynthetic pathway. β-carotene is the most abundant carotenenes in the leaves since they were the primary free radical scavenger in the photosynthesis whereas α-carotene was present as minor accessory pigment in a number of plant species only[34].




Fig. 1: Difference in the structures of α-carotene and β-carotene.

As this paperwork mainly discussed on pro-vitamin A content of the selected *ulam* species, hence the related retinol equivalent activity of respective samples was also included here for further evaluation on their relevancy as natural sources for vitamin A pre-cursors. The total retinol equivalent (RE) content for every sample was tabulated in Table 3. The total RE content was calculated in accordance to NRC[23,35]. Attentively, *Beluntas* ranked at the top to possess 656.59 µg RE per g DW, followed by *Cekur Manis*, *Pegaga* and *Daun Selom* with 592.45 µg, 576.50 µg and 514.17 µg RE/g DW respectively.

Table 3: Total retinol equivalent (RE) activity in selected 22 *ulam* species

Samples	Total RE	Samples	Total RE
<i>Salad</i>	90.37	<i>Daun Bawang</i>	344.48
<i>Kemangi</i>	130.57	<i>Selasih</i>	368.52
<i>Mas Cotek</i>	145.77	<i>Daun Kucai</i>	369.27
<i>Tenggek Burung</i>	209.97	<i>Sirih</i>	377.42
<i>Kaduk</i>	211.43	<i>Daun Sup</i>	462.21
<i>Beka</i>	219.34	<i>Ulam Raja</i>	464.54
<i>Daun Kari</i>	281.69	<i>Daun Selom</i>	481.34
<i>Sawi</i>	284.36	<i>Daun Kesum</i>	514.17
<i>Gajus</i>	289.69	<i>Pegaga</i>	576.50
<i>Hempedu Bumi</i>	294.36	<i>Cekur Manis</i>	592.45
<i>Jarum Tujuh Bilah</i>	310.60	<i>Beluntas</i>	656.59

Data represented in microgram per gram dry weight.

According to Ministry of Health (MOH), Malaysia, the recommended nutrient intakes (RNI) for vitamin A lies within range 375 to 850 µg RE per day (about 2 to 5 mg β-carotene per day). However, these values become specific based on gender, age and stage of life⁵. From the analysis done in this study, RE values from eight of the 22 samples established to be within the range as suggested by both MOH and FAO/WHO[5,36]. There have been reviews on several plants to be used as interventional food in order to overcome VAD issue such as orange-fleshed carrots and orange-fleshed sweet potatoes[37,38]. As for comparison, analysis for orange-fleshed carrots was also carried out to check for their RE content which valued at 99.64 µg RE/g DW. Thereby, from the results obtained earlier, we have found that 21 out of the 22 selected *ulam* species are having better RE values than carrots ranged from 130.57 to 656.59 µg RE/g DW. The individual pro-vitamin A profiles are well-established here and thus providing an update on the nutritional values of our traditional vegetables (*ulam*). Further investigation need to be taken into account before the *ulam* species could be used for local food intervention program to eliminate VAD as a public health concern.

CONCLUSION

The retinol equivalent (RE) values in every sample are mainly correlated to their respective concentration of β-carotene. It was

found that 21 of the analysed *ulam* species are having better RE activity in comparison to a well-known source of pro-vitamin A (orange fleshed carrots), hence could be further evaluated for local food intervention program to combat VAD problems.

REFERENCES

- World Health Organization (WHO). Global prevalence of vitamin A deficiency in populations at risk 1995–2005. WHO Global Database on Vitamin A Deficiency. Geneva: WHO; 2009.
- United Nations Children's Fund (UNICEF). Vitamin A: A decade of progress. New York: UNICEF; 2007.
- Khor GL. Micronutrient deficiency and its alleviation: The case of Malaysia. Asia Pacific Journal of Clinical Nutrition 2002, 11 (Suppl.): S377-S381.
- Institute of Medicine. Vitamin A. In: Dietary reference intakes for vitamin A, vitamin K, arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium, and zinc. Food and Nutrition Board, Institute of Medicine. Washington, D.C. National Academy Press; 2001: 82–161.
- Ministry of Health. Vitamin A. In: Recommended nutrient intakes for Malaysia. National Coordinating Committee on Food and Nutrition, Ministry of Health Malaysia. Putrajaya; 2005: 111-120.

6. Devadas RP, Saroja S. Availability of Fe and β -carotene from *Amaranthus* to children. In: Emmaus P, Editor. Proceedings of the second *Amaranthus* conference, Rodala Press Inc.; 1980: 15-21.
7. Tee ES, Lim CL. Carotenoids composition and content of Malaysian vegetables and fruits by the AOAC and HPLC methods. *Food Chemistry* 1991; 41: 309-339.
8. Rodriguez-Amaya DB. Important Food Sources of Provitamins A. Carotenoids and food preparation: The retention of provitamin A carotenoids in prepared, processed, and stored foods. OMNI Press: Washington; 1997.
9. Liu Y, Perera CO, Suresh V. Comparison of three chosen vegetables with others from South East Asia for their lutein and zeaxanthin content. *Food Chemistry* 2007; 101: 1533-1539.
10. Raju M, Varakumar S, Lakshminarayana R, Krishnakantha TP, Baskaran V. Carotenoid composition and vitamin A activity of medicinally important green leafy vegetables. *Food Chemistry* 2007; 101: 1598-1605.
11. Kopsell DA, Kopsell DE. Carotenoids in vegetables: Biosynthesis, occurrence, impacts on human health, and potential for manipulation. In: Watson RR, Preedy VR, Editors. *Bioactive foods in promoting health: Fruits and vegetables*. Elsevier Inc.: Boston; 2010: 645-662.
12. Seddon JM, Ajani UA, Sperduto RD, Hiller R, Blair N, Burton TC *et al.* Dietary carotenoids, vitamins A, C and E, and advanced age-related macular degeneration. *Journal of American Medical Association* 1994; 272 (18): 1413-1420.
13. Landrum JT, Bone RA, Kilburn MD. The macular pigment: A possible role in protection from age-related macular degeneration. *Advances in Pharmacology* 1997; 38: 537-556.
14. Cunningham FX Jr, Gantt E. Genes and enzymes of carotenoid biosynthesis in plants. *Annual Review of Plant Physiology and Plant Molecular Biology* 1998; 49: 557-583.
15. Paiva ARS, Russell RM. β -carotene and other carotenoids as antioxidants. *Journal of the American College of Nutrition* 1999; 18 (5): 426-433.
16. Garcia-Closas R, Berenguer A, Tormo MJ, Sanchez MJ, Quiros JR, Navarro C *et al.* Dietary sources of vitamin C, vitamin E and specific carotenoids in Spain. *British Journal of Nutrition* 2004; 91: 1005-1011.
17. Saidin I. *Sayuran tradisional, ulam dan penyedap rasa*. Penerbit Universiti Kebangsaan Malaysia: Bangi; 2000.
18. Ong HC. *Sayuran: Khasiat Makanan dan Ubatan*. Utusan Publications and Distributors: Kuala Lumpur, Malaysia; 2003.
19. Ong HC. *Sayuran 2: Khasiat Makanan dan Ubatan*. Utusan Publications and Distributors: Kuala Lumpur, Malaysia; 2011.
20. Mohd Zaifuddin FA, Mohd Hassan N, Othman R. Identification of carotenoid composition in selected '*ulam*' or traditional vegetables of Malaysia. *International Food Research Journal* 2012; 19(2): 527-530.
21. Othman R. *Biochemistry and genetics of carotenoid composition in potato tubers*. Christchurch, New Zealand: Lincoln University, PhD thesis; 2009.
22. Rodriguez-Amaya DB, Kimura M. *Harvest Plus handbook for carotenoid analysis*. Harvest Plus: Washington DC; 2004.
23. National Research Council. *Recommended Dietary Allowances*. National Academy Press: Washington DC; 1989.
24. Andarwulan N, Kurniasih D, Apriady RA, Rahmat H, Roto AV, Bolling BW. Polyphenols, carotenoids and ascorbic acid in underutilized medicinal vegetables. *Journal of Functional Foods* 2012; 4(1): 339-347.
25. Speek AJ, Speek-Saichua S, Schreurs WHP. Total carotenoid and β -carotene contents of Thai vegetables and the effect of processing. *Food Chemistry* 1988; 27 (4): 245-257.
26. Wills RBH, Ranga A. Determination of carotenoids in Chinese vegetables. *Food Chemistry* 1996; 56 (4): 451-455.
27. Burns J, Fraser PD, Bramley PM. Identification and quantification of carotenoids, tocopherols and chlorophylls in commonly consumed fruits and vegetables. *Phytochemistry* 2003; 62: 939-947.
28. Mohd Salleh MR. *Penyarian zeaxanthin dari sayur-sayuran tempatan: Perbandingan Kaedah penyarian dan kesan parameter proses terhadap hasil sarianzeaxanthin*. Skudai, Malaysia: University Technology Malaysia, MSc thesis; 2005.
29. Lakshminarayana R, Raju M, Krishnakantha TP, Baskaran V. Lutein and zeaxanthin in leafy greens and their bioavailability: Olive oil influences the absorption of dietary lutein and its accumulation in adult rats. *Journal of Agricultural and Food Chemistry* 2007; 55 (15): 6395-6400.
30. Murillo E, Melendez-Martinez AJ, Portugal, F. Screening of vegetables and fruits from Panama for rich sources of lutein and zeaxanthin. *Food Chemistry* 2010; 122 (1): 167-172.
31. Khoo HE, Prasad KN, Kong KW, Jiang Y, Ismail A. Carotenoids and their isomers: Color pigments in fruits and vegetables. *Molecules* 2011; 16 (2): 1710-1738.
32. Mamatha BS, Sangeetha RK, Baskaran V. Provitamin-A and xanthophyll carotenoids in vegetables and food grains of nutritional and medicinal importance. *International Journal of Food Science and Technology* 2011; 46: 315-323.
33. Aruna G, Baskaran V. Comparative study on the levels of carotenoids lutein, zeaxanthin and β -carotene in Indian spices of nutritional and medicinal importance. *Food Chemistry* 2010; 123 (2): 404-409.
34. Demmig-Adams B, Gilmore AM, Adams WW. In vivo functions of carotenoids in higher plants. *Journal of the Federation of American Societies for Experimental Biology* 1996; 10: 403-412.
35. Eitenmiller RR, Ye L, Landen WO. Vitamin A and carotenoids. In: *Vitamin analysis for the health and food sciences*. 2nd edition. Florida: CRC Press; 2008: 3-81.
36. FAO/WHO. *Vitamin A*. In: *Human vitamin and mineral Requirements*. Report of a joint FAO/WHO expert consultation. FAO: Rome; 2002: 87-107.
37. Arscott SA, Tanumihardjo SA. Carrots of many colors provide basic nutrition and bioavailable phytochemicals acting as a functional food. *Comprehensive Review in Food Science and Food Safety* 2010; 9: 223-239.
38. Burri BJ. Evaluating sweet potato as an intervention food to prevent vitamin A deficiency. *Comprehensive Reviews in Food Science and Food Safety* 2011; 10 (2): 118-130.